

correction dose

This is the insulin dose you take for high blood glucose during the day and at bedtime.

Meal time: 1 unit per 5mg/dL over 150

Bedtime: 1 unit per 5mg/dL over 155

blood glucose	daytime	bedtime
Under 150	0	0
150-155	1	0
156-160	2	1
161-165	3	2
166-170	4	3
171-175	5	4
176-180	6	5
181-185	7	6
186-190	8	7
191-195	9	8
196-200	10	9
201-205	11	10
206-210	12	11
211-215	13	12
216-220	14	13
221-225	15	14
226-230	16	15

continued on back

Only give correction dose if it has been **more than 2-3 hours** since the last dose of fast-acting insulin.

blood glucose	daytime	bedtime
231-235	17	16
236-240	18	17
241-245	19	18
246-250	20	19
251-255	21	20
256-260	22	21
261-265	23	22
266-270	24	23
271-275	25	24
276-280	26	25
281-285	27	26
286-290	28	27
291-295	29	28
296-300	30	29
301-305	31	30
306-310	32	31
311-315	33	32
316-320	34	33
321-325	35	34
326-330	36	35
331-335	37	36
336-340	38	37
341-345	39	38
346-350	40	39
351-355	41	40
356-360	42	41

childrensdayton.org/diabetesresources

